

Gyn Cancer & Pelvic Surgery ^{LLC}

Practice limited to Gynecologic Oncology, Reconstructive Pelvic Surgery and Consultative Gynecology

THAD R. DENEHY, M.D., FACOG, FACS

BOARD CERTIFIED IN GYN ONCOLOGY & OB/GYN

MEDICATIONS TO DISCONTINUE PRIOR TO SURGERY

*Please stop/discontinue taking
any of the following medications
7 days prior to surgery:*

- Anti-Platelet Agents (Plavix)
- Over-the-Counter Medications
- Herbal Remedies
- Nutritional Supplements
- Ibuprofen
- Motrin
- Advil
- Bufferin
- Excedrin
- Aleve
- Ecotrin
- Anacin
- Pletal
- Alka-Seltzer
- Arthritis Formulas
- Vitamin E
- Garlic Capsules
- Ginko Biloba
- Multivitamin Formulas

*Please stop/discontinue taking
any of the following
medications prior to surgery:*

- Coumadin (Warfarin) - 4 days
- Lovenox - as directed by your
physician
- Heparin - as directed by your
physician

Please discontinue any birth control pills
1 month before scheduled procedure
(unless otherwise specified by your
specialist).

If you take high blood pressure, heart,
diabetic, or any other medications consult
with your **physician that prescribed** these
medications whether or not you may take it
the morning of your procedure.

Remember to drink plenty of **liquids** the
day before your procedure.